

## NP-HARD and NP-COMPLETE Problems

- NP-HARD and NP-COMPLETE Problems
- A first NP-COMPLETE Problem: CIRCUIT-SAT( $C$ )
- The Cook-Levin Theorem: SAT is NP-COMPLETE
- NP-COMPLETE Problems from known Reductions
- NP-COMPLETE ness of DIR-HAM-CYCLE and HAM-CYCLE
- TSP is NP-COMPLETE
- SUBSET-SUM is NP-COMPLETE
- PARTITION is NP-COMPLETE

IMDAD ULLAH KHAN

## Proving NP-COMPLETE Problems

---

A problem  $X$  is **NP-COMPLETE**, if

- 1  $X \in \text{NP}$
- 2  $\forall Y \in \text{NP } Y \leq_p X$

How to prove a problem NP-COMPLETE?

To prove  $X$  to be NP-COMPLETE

- 1 Prove  $X \in \text{NP}$
- 2 Reduce some known NP-COMPLETE problem  $Z$  to  $X$

Again! Reduce a known NP-COMPLETE problem to  $X$

▷ **Not the other way round. A very common mistake!**

## A first NP-COMPLETE Problem

---

### Theorem (The Cook-Levin theorem)

$SAT(f)$  is NP-COMPLETE

- Proved by Stephen Cook (1971) and earlier by Leonid Levin (but became known later)
- Levin proved six NP-COMPLETE problems (in addition to other results)
- We prove this by reducing  $CIRCUIT-SAT(C)$  problem to  $SAT(f)$  problem

## A first NP-COMPLETE Problem

---

To prove  $X$  NP-COMPLETE, reduce an NP-COMPLETE problem  $Z$  to  $X$

Where to begin? we need a first NP-COMPLETE Problem

Theorem (The Cook-Levin theorem)

$SAT(f)$  is NP-COMPLETE

- Proved by Stephen Cook (1971) and earlier by Leonid Levin (but became known later)
- Levin proved six NP-COMPLETE problems (in addition to other results)

We prove the theorem by reducing  $CIRCUIT-SAT(C)$  problem to  $SAT(f)$  problem

## The Cook-Levin theorem

### Theorem (The Cook-Levin theorem)

$SAT(f)$  is NP-COMPLETE

We already showed that SAT is polynomial time verifiable

$SAT \in NP$

Now we prove that

$CIRCUIT-SAT(C) \leq_p SAT(f)$

This proves that SAT is NP-HARD and completes the proof

- Suppose  $\mathcal{A}$  is an algorithm to decide  $SAT(f)$
- Given an instance  $C$  of the  $CIRCUIT-SAT(C)$  problem
- In polynomial time we transform  $C$  into an equivalent CNF formula  $f$
- Make a call  $\mathcal{A}(f)$  to decide whether or not  $CIRCUIT-SAT(C) = \mathbf{Yes}$

# The Cook-Levin theorem

$$\text{CIRCUIT-SAT}(C) \leq_p \text{SAT}(f)$$

Make a variable for each input wire and output of each gate of the circuit  $C$

For each not gate make **equi-satisfiable clauses**





- These clauses are satisfied iff  $x_j = \bar{x}_i$

For each and gate make **equi-satisfiable clauses**

- These clauses are satisfied iff  $x_k = x_i \wedge x_j$

For each or gate make **equi-satisfiable clauses**

- These clauses are satisfied iff  $x_k = x_i \vee x_j$



## The Cook-Levin theorem

---

$$\text{CIRCUIT-SAT}(C) \leq_p \text{SAT}(f)$$

- Easy to verify that the gates and corresponding formula are equisatisfiable
- The output gate value is encoded with a clause containing the corresponding variable
- The final formula  $f$  is a grand conjunction of all the clauses made for each gate and output of the circuit  $C$

$f$  is equisatisfiable with the  $C$


▷ i.e.  $\text{CIRCUIT-SAT}(C) = \text{Yes}$  if and only if  $\mathcal{A}(f) = \text{Yes}$

The reduction takes polynomial time, requires one traversal of the DAG, constant time per gate

## Implied NP-COMPLETE Problems

From known reductions, the following problems are NP-COMPLETE

- $SAT \leq_p 3-SAT$
- $3-SAT \leq_p IND-SET$
- $IND-SET \leq_p CLIQUE$
- $IND-SET \leq_p VERTEX-COVER$
- $VERTEX-COVER \leq_p SET-COVER$
- $IND-SET \leq_p SET-PACKING$



We show a few more reductions to prove problems to be NP-COMPLETE



We showed DIR-HAM-CYCLE to be in NP for NP-HARDNESS we prove

$$3\text{-SAT}(f) \leq_p \text{DIR-HAM-CYCLE}(G)$$

Let  $f$  be an instance of 3-SAT on  $n$  variables and  $m$  clauses

Let  $x_1, \dots, x_n$  be the variables and  $C_1, \dots, C_m$  be the clauses of  $f$


Construct a digraph  $G$  that has a Hamiltonian cycle iff  $f$  is satisfiable

- 1 In  $G$  there will be  $2^n$  sub-Hamiltonian cycles corresponding to the  $2^n$  possible assignments to variables  $x_1, \dots, x_n$
- 2 We introduce a structure for each clause such that these sub-Hamiltonian cycles can be combined if and only if all clauses are satisfiable


## DIR-HAM-CYCLE is NP-COMPLETE

$$3\text{-SAT}(f) \leq_p \text{DIR-HAM-CYCLE}(G)$$

For each  $x_i$  make a sequence of  $3(m+1)$  bidirectionally adjacent vertices




- $x_i = 1 \implies$  traverse this gadget from  $L_i$  to  $R_i$  and vice-versa
- $(x_i, x_{i+1}) = (1, 0) \implies$  traverse from  $L_i \rightarrow R_i \rightarrow R_{i+1} \rightarrow L_{i+1}$
- $(x_i, x_{i+1}) = (0, 0) \implies$  traverse from  $R_i \rightarrow L_i \rightarrow R_{i+1} \rightarrow L_{i+1}$



## DIR-HAM-CYCLE is NP-COMPLETE

$$3\text{-SAT}(f) \leq_p \text{DIR-HAM-CYCLE}(G)$$


Make nodes  $s$  and  $t$  and combine all the gadgets as follows



## DIR-HAM-CYCLE is NP-COMPLETE

$$3\text{-SAT}(f) \leq_p \text{DIR-HAM-CYCLE}(G)$$

- $2^n$  Ham cycles traversing each gadget in either direction
- These correspond to the  $2^n$  possible assignments to the  $n$  variables
- **Make a Hamiltonian cycle exist iff there is a satisfying assignment**
- Have to incorporate clauses. Make nodes for each clause
- If a variable satisfy a clause, traverse it by a detour from that gadget



## DIR-HAM-CYCLE is NP-COMplete

---

$$3\text{-SAT}(f) \leq_p \text{DIR-HAM-CYCLE}(G)$$

Given  $f$ , make  $G$  as described above

$G$  has a directed Hamiltonian cycle iff  $f$  is satisfiable


The construction takes polynomial time (about  $O(nm)$ )

## DIR-HAM-PATH is NP-COMPLETE

$$\text{DIR-HAM-CYCLE}(G) \leq_p \text{DIR-HAM-PATH}(G')$$

Let  $G = (V, E)$  be an instance of the  $\text{DIR-HAM-CYCLE}(G)$  problem

- For any arbitrary  $v \in V$ , make  $G'$  on  $V(G) \setminus \{v\} \cup \{v_{in}, v_{out}\}$ 
  - ▷ i.e. remove  $v$  and add two new vertices  $v_{in}$  and  $v_{out}$
- $v_{in}$  has all incoming edges of  $v$  directed to it from in-neighbors of  $v$
- $v_{out}$  has all outgoing edges of  $v$  directed from it to out-neighbors of  $v$



$G$  has a directed Hamiltonian cycle iff  $G'$  has a directed Hamiltonian path


## HAM-CYCLE is NP-COMPLETE

We proved its polynomial time verifiability earlier, now we show that

$$\text{DIR-HAM-CYCLE}(G) \leq_p \text{HAM-CYCLE}(G')$$

Let  $G = (V, E)$  be an instance of the  $\text{DIR-HAM-CYCLE}(G)$ .  $|V| = n$ ,  $|E| = m$

- Make an undirected graph  $G' = (V', E')$ ,  $|V'| = 3n$  and  $|E'| = m + 2n$
- Split every vertex  $v \in V$  into three vertices  $v_{in}$ ,  $v_{md}$ ,  $v_{ot}$  and add to  $V'$
- Add edges  $(v_{in}, v_{md})$  and  $(v_{md}, v_{ot})$  in  $E'$
- For each directed edge  $(x, y) \in E$ , make the edge  $(x_{ot}, y_{in})$  in  $E'$




$G$  has a dir-Ham cycle iff  $G'$  has an (undirected) Hamiltonian cycle

## TSP is NP-COMPLETE


$$\text{HAM-CYCLE}(G) \leq_p \text{TSP}(G', k)$$

- $\text{TSP}(G', k)$  requires weighted graph and a number  $k$
- Given an instance  $G = (V, E)$  of  $\text{HAM-CYCLE}(G)$ ,  $|V| = n$
- Make a complete graph on  $n$  vertices  $G'$  with weights as follows


$$w(v_i, v_j) = \begin{cases} 1 & \text{if } (v_i, v_j) \in E(G) \\ 2 & \text{else} \end{cases}$$




Hamiltonian cycle  
in  $G$  shown in blue



TSP tour in  $G'$  of  
of length shown in blue



No Hamiltonian  
cycle in  $G$



No TSP tour of  
length 5 in  $G'$

$G$  has a Hamiltonian cycle iff  $G'$  has a TSP tour of length  $k = n$



### SUBSET-SUM is NP-COMPLETE

- Given a set  $U = \{a_1, a_2, \dots, a_n\}$  of integers
- A weight function  $w : U \rightarrow \mathbb{Z}^+$ , and a positive integer  $C$
- The SUBSET-SUM( $U, w, C$ ) problem: Is there a  $S \subset U$  with  $\sum_{a_i \in S} w_i = C$ ?
- If  $w_i$ 's and  $C$  are given in unary encoding
  - then  $O(nC)$  dynamic programming solution is a polynomial time
- But this is exponential in size of input if  $C$  is provided in binary (or decimal)

We prove that

$$3\text{-SAT}(f) \leq_p \text{SUBSET-SUM}(\bullet, \bullet, \bullet)$$

## SUBSET-SUM is NP-COMPLETE

$$3\text{-SAT}(f) \leq_p \text{SUBSET-SUM}(\bullet, \bullet, \bullet)$$

- Given an instance  $f$  of  $3\text{-SAT}(f)$  with  $n$  variables and  $m$  clauses
- Construct  $2n + 2m$  weights: 2 objects for each variable and each clause
- Each is a  $n + m$ -digits integer (a digit for each variable and each clause)
- The weight for literal  $x_i$  and  $\bar{x}_i$  have digit 1 corresponding to the variable  $x_i$
- The digit for clause  $C_j$  is 1 if the literal appears in clause  $C_j$

	$x_1$	$x_2$	$x_3$							$x_{n-1}$	$x_n$	$C_1$	$C_2$	$C_3$		$C_{m-1}$	$C_m$
$x_1$	1								...			1	1				
$\bar{x}_1$	1								...					1			
$x_2$		1							...			1	1			1	1
$\bar{x}_2$		1							...					1			
$x_3$			1						...								1
$\bar{x}_3$			1						...			1	1	1			
$\vdots$			$\vdots$									$\vdots$	$\vdots$	$\vdots$			$\vdots$

$$C_1 = (x_1 \vee x_2 \vee \bar{x}_3), C_2 = (x_1 \vee x_2 \vee \bar{x}_3), C_3 = (\bar{x}_1 \vee \bar{x}_2 \vee \bar{x}_3)$$

$$C_{m-1} = (x_2 \vee x_8 \vee x_9), C_m = (x_2 \vee x_3 \vee x_5)$$

# SUBSET-SUM is NP-COMPLETE

$$3\text{-SAT}(f) \leq_p \text{SUBSET-SUM}(\bullet, \bullet, \bullet)$$

- Remaining  $2m$  weights set so as last sum of digits at each position from  $n + 1$  to  $n + m$  is 5 ▷ details in notes

	$x_1$	$x_2$	$x_3$	...	$x_{n-1}$	$x_n$	$C_1$	$C_2$	$C_3$	...	$C_{m-1}$	$C_m$
$x_1$	1							1	1			
$\bar{x}_1$	1								1			
$x_2$		1						1	1			1
$\bar{x}_2$		1							1			
$x_3$			1									1
$\bar{x}_3$			1						1	1	1	
$\vdots$												
$x_n$												
$\bar{x}_n$												
								1				
								1				
									1			
										1		
												1
												1

# SUBSET-SUM is NP-COMPLETE

$$3\text{-SAT}(f) \leq_p \text{SUBSET-SUM}(\bullet, \bullet, \bullet)$$

	$x_1$	$x_2$	$x_3$	...	$x_{n-1}$	$x_n$	$C_1$	$C_2$	$C_3$	...	$C_{m-1}$	$C_m$
$x_1$	1							1	1			
$x_2$	1								1	1		
$x_3$		1						1	1			1
$x_4$			1							1		
$x_5$				1								1
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$x_n$												
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$x_{n+m}$							1					
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	1					
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$		1				
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$			1			
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$				1		
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$					1	
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$						1

The SUBSET-SUM instance with  $2n + 2m$  weights as shown above and

$$C = \underbrace{111\dots, 11}_n \underbrace{333\dots 33}_m \text{ is Yes if and only if } f \text{ is satisfiable}$$

## PARTITION is NP-COMPLETE

$$\text{SUBSET-SUM}(U, w, C) \leq_p \text{PARTITION}(U', k)$$

- Let  $U' = \{w_1, w_2, \dots, w_n, w_{n+1}, w_{n+2}\}$
- $w_{n+1} = 2\left[\sum_{i=1}^n w_i\right] - C$  and  $w_{n+2} = \left[\sum_{i=1}^n w_i\right] + C$

$\text{SUBSET-SUM}(U, w, C) = \mathbf{Yes}$  iff  $\text{PARTITION}(U', \mathbf{0}) = \mathbf{Yes}$  (balanced)

- $\sum_{x \in U'} x = \sum_{a_i \in U} w_i + 2\left[\underbrace{\sum_{i=1}^n w_i}_{w_{n+1}}\right] - C + \left[\underbrace{\sum_{i=1}^n w_i}_{w_{n+2}}\right] + C = 4 \sum_{a_i \in U} w_i$
- Let  $P_1$  and  $P_2$  be a balanced bipartition of  $U'$
- Both  $w_{n+1}$  and  $w_{n+2}$  cannot be in the same part, assume  $w_{n+1} \in P_1$
- Both  $P_1$  and  $P_2$  cannot contain only one element, so  $\sum_{x \in P_1 \setminus \{w_{n+1}\}} w_x = C$

 $P_1$ 
 $P_2$ 

$$w_{n+1} = 2 \sum_i w_i - C \quad C$$

$$w_{n+2} = \sum_i w_i + C \quad \sum_i w_i - C$$

## NP-COMPLETE Problems

21 problems were shown to be NP-COMPLETE in a seminal paper: Richard Karp (1972), "Reducibility Among Combinatorial Problems"




FIGURE 1 - Complete Problems

RICHARD M. KARP