

Approximation Algorithms

- Approximation Algorithms for Optimization Problems: Types
- Absolute Approximation Algorithms
- Inapproximability by Absolute Approximate Algorithms
- Relative Approximation Algorithm
- Inapproximability by Relative Approximate Algorithms
- Polynomial Time Approximation Schemes
- Fully Polynomial Time Approximation Schemes

Absolute Approximation Algorithms

Given an optimization problem P with value function f on solution space

An algorithm A is called **absolute approximation** algorithm if there is a constant k such that for any instance I

$$|f(A(I)) - f(\text{OPT}(I))| \leq k$$

- For a minimization problem this means $f(A(I)) \leq f(\text{OPT}(I)) + k$
- For a maximization problem this means $f(A(I)) \geq f(\text{OPT}(I)) - k$

How to Design an Approximation Algorithm?

Any approximation algorithm involves 2 main ingredients:


1 First step is to design a good algorithm A

For approximation guarantee on $A(I)$ we need the value of the optimal solution $f(\text{OPT}(I))$

▷ How to find it? almost equally difficult (version of the problem)


2 We find a good lower or upper bound on $f(\text{OPT}(I))$

3 Compare $f(A(I))$ with the bound on $f(\text{OPT}(I))$




Graph Coloring


A **graph (vertex) coloring** is to assign a color to each vertex such that no two adjacent vertices get the same color




A graph G on 8 vertices



A coloring with 6 colors



A coloring with 8 colors



A coloring with (optimal) 3 colors

[**COLORING(G)** problem:] Color G with minimum number of colors, $\chi(G)$

Graph Coloring (Optimization) Problem


The graph coloring (optimization) problem

- \mathcal{I} : Graphs
- $S(I)$: An assignment of colors to vertices of input graph, such that no two adjacent vertices have the same color (feasibility)
- $f : S(I) \rightarrow \mathbb{Z}^+$
 - For $s \in S(I)$, $f(s)$ is number of colors used in the coloring s
- $\chi(G)$: the minimum number of colors needed to color G
 - $\chi(G) = f(\text{OPT}(I))$

Planar Graphs

Planar Graphs

A graph is planar if it can be drawn in the plane without any edge crossing




Four different drawings of the same graph, K_4

Planar Graphs

Planar Graphs

A graph is planar if it can be drawn in the plane without any edge crossing

Just because in a drawing of G edges cross does not mean G is not planar




Planar Graphs

Planar Graphs

A graph is planar if it can be drawn in the plane without any edge crossing

To prove planarity, move the vertices around, redraw the edges without crossing (sometimes in a very indirect fashion)




Planar Graphs

Planar Graphs


A graph is planar if it can be drawn in the plane without any edge crossing

To prove planarity, move the vertices around, redraw the edges without crossing (sometimes in a very indirect fashion)

Even harder to prove non-planarity



K_5



$K_{3,3}$

Planar Graphs: A characterization

A graph is planar if it can be drawn in the plane without any edge crossing


We will find some invariants that planar graphs satisfy

To prove non-planarity of a graph G , we will show that G doesn't satisfy that invariant

Planar Graphs: A characterization

A plane drawing of a planar graph divides the plane into regions or faces, one of them the outer face

A **region (or face)** is a part of the plane disconnected from other parts by the edges




Planar Graphs: A characterization



Euler's Formula: The number of faces of a connected planar graph is invariant of its drawing and is given by

$$f = e - v + 2 \quad f = |\text{faces}|, \quad e = |E|, \quad v = |V|$$

Verify it for the following planar graphs



outer face



Planar Graphs: A characterization

Euler's Formula: The number of faces of a connected planar graph is invariant of its drawing and is given by

$$f = e - v + 2 \quad f = |\text{faces}|, \quad e = |E|, \quad v = |V|$$

Face-Edge Handshaking Lemma

Let G be a planar graph, let R be its regions, then

$$2e = \sum_{F \in R} \text{deg}(F)$$

If G is a connected planar graph with $v \geq 3$, then $e \leq 3v - 6$

An immediate corollary from this using the Handshaking Lemma is

Every planar graph has a vertex with degree at most 5

Coloring Planar Graphs

The **PLANAR-GRAPH-COLORING(G)** problems: Given a planar graph G , color it with minimum colors

6-Coloring Planar Graphs

Every planar graph has a vertex with degree at most 5

Using this lemma we give a recursive 6-coloring algorithm

▷ Can apply the algorithm to components of disconnected graphs

Algorithm 6-COLOR($G, C = \{c_1, \dots, c_6\}$)


Let $v \in V$ such that $\deg(v) \leq 5$

6-COLOR($G - v, C = \{c_1, \dots, c_6\}$)

Let $C' \subset C$ be the set of colors used for $N(v)$

▷ $|C'| \leq 5$

Color v with a color in $C \setminus C'$



▷ Clearly polynomial time

6-Coloring Planar Graphs


Every planar graph has a vertex with degree at most 5

Using the lemma we give a recursive 6-coloring algorithm

▷ Can apply the algorithm to components of disconnected graphs

Can be implemented with no recursion or modification to adjacency list

- 1 Order vertices so no vertex has more than 5 neighbors preceding it
- 2 Greedily color vertices from left to right using the scheme in figure



▷ Clearly polynomial time

3-absolute approximation for Planar Graph Coloring

Theorem: The decision problem of planar graph 3-Coloring is NP-COMPLETE

Use the fact that: If G is bipartite, then it is 2-colorable

Algorithm APPROX-PLANAR-COLOR(G)

if G is bipartite **then**

▷ Easy to check with a BFS

Color G with the obvious 2-coloring

else

6-COLOR($G, C = \{c_1, \dots, c_6\}$)

APPROX-PLANAR-COLOR is a 3-absolute approximate algorithm

- Non-bipartite graphs require ≥ 3 colors ($f(\text{OPT}(G)) \geq 3$) ▷ **(LB)**
- We use at most 6 colors ▷ $f(\text{APPROX-PLANAR-COLOR}(G)) \leq 6$
- The statement follows


2 and 1-absolute approximation for Planar Graph Coloring

- A slightly complicated algorithm colors planar graphs with 5 colors

That algorithm due to Kempe, is a 2-absolute approximate algorithm

- Appel and Haken (1976) gave a complicated proof that planar graphs can be colored with 4 colors

That “algorithm” is a 1-absolute approximate algorithm



UIUC stamp in honor of the 4-Color theorem